


OENOBrett®

Chitosan and enzyme based preparation for the control of *Brettanomyces* spoilage.
 Qualified for the elaboration of products for direct human consumption in the field of the regulated use in Oenology. In accordance with the regulation (EC) n° 2019/934.




Chitosan is extracted from the fungus *Aspergillus niger*. It is therefore of non-animal origin and is also allergen-free. Its antagonism towards yeasts has been scientifically proven.

SPECIFICATIONS AND OENOLOGICAL PROPERTIES

OENOBrett® is a specific combination of a natural polysaccharide: chitosan and a pectinase / glucanase enzymatic preparation that facilitates the lysis and the elimination of *Brettanomyces* yeasts.


EXPERIMENTAL RESULTS

- Trials on Cabernet Sauvignon 2011 (free SO₂= 21 mg/L, total SO₂= 85 mg/L, pH=3.85, temperature = 15°C, OENOBrett® was added to the wine (10 g/hL) while a control received 20 mg/L SO₂. Additions were made simultaneously and wines were racked at the same time.



The synergy of the chitosan and enzyme preparation ensures the efficiency of **œNOBrett®** regardless of:

- The population level, the physiological state and the *Brettanomyces* strain.
- The colloidal state of the wine.



PHYSICAL CHARACTERISTICS

Aspect thin powder

Colour clear beige

CHEMICAL AND MICROBIOLOGICAL ANALYSIS

Acetylation degree (mol %) 0 - 30
 Humidity (%) < 10
 Coliformes (CFU/g) < 60
E. coli (/25 g) none

Salmonella (/25 g) none
 Lead (ppm) < 3.5
 Arsenic (ppm) < 2.5
 Mercury (ppm) < 0.4

PROTOCOL FOR USE

OENOLOGICAL CONDITIONS

- Treatment must be after the completion of both AF and MLF.
- Adjust wine SO₂ in accordance to good oenological practices.

DOSAGE

Recommended dosage: 10 g/hL (100 ppm).
 EU Regulation: Maximum legal dosage: 25 g/hL (250 ppm).

IMPLEMENTATION

- Dissolve in ten times its weight in water or wine – Use a clean and inert container.
- Introduce the preparation into the wine, homogenise properly (maintain a stable wine temperature above 10°C throughout the treatment).
- Possible racking after 8 days of treatment.

STORAGE RECOMMENDATION

- Store above ground level in a dry area not liable to impart odours. Ensuring stock is kept at a moderate temperature, in its original, unopened packaging.
- Optimal date of use: 3 years.

PACKAGING

250 g and 2.5 kg boxes.
 23 g doses, 460 g boxes (20 x 23 g).

